

Clinical Study Of Enteropathogenic Which Causes Diarrhea Among ٠- ١٥ Years Age Group In Baghdad City

Khalida Jhalil Ibraheem

Instructor, M.Sc. of Microbiology
Dept. of Sciences , College of Basic Education
Al-Mustansiriyah University

Abstract

Diarrheal diseases are a leading cause of mortality among children in developing countries. This study is designed to investigate the bacterial, viral and parasitic etiology and related clinical and epidemiological factors in children among ٠-١٥ years age group with diarrhea in Baghdad city. A total of ٢٠٠ sample of the stool obtained from children with diarrhea visiting child's Central Teaching Hospital and Alkadhmiya Hospital for children in the period between April , ٢٠١٦ to October , ٢٠١٦ . Stool sample were examined by using standard culture and identification methods for bacteria , and by microscopically using normal saline and iodine staining wet mounts for identification of parasites and by Rapid Diagnostic Kit for detection of rotavirus. From summation of ٢٠٠ cases in the study group , ١٤٤ (٧٢%) infected by single etiology and ٥٦ (٢٨%) infected by mixed etiology . The highest infection rate recorded in the bacterial group (٥١,٥%) followed by parasitic group (٣٧,٥%) and viral group (١١%) . According to the age categories , it was observed that maximum bacterial infection rate at age group (< ١ years) were (٤٦%), while at age group (> ٥ years) were (٥٢%) for maximum parasitic infection and viral infection recorded the highest rate in the age group (١-٥ years) by ٧٣% . The result of the study showed that the predominant bacteria were *Escherichia coli* with ٢٠,٥% followed by *Shigella sp.* , *Salmonella sp.* and *Klebsiella sp.* with ١٤,٥% , ١٠% and ٦,٥% respectively . *Entamoeba histolytica* recorded highest parasitic infection rate ١٩,٥% followed by *Giardia lamblia* ١٢,٥% and *Blantidium coli* ٥,٥% . The percentage of samples infected with rotavirus was ١١% compared with the total number of samples studied .

Introduction

Clinical Study Of Enteropathogenic Which Causes Diarrhea Among ٠-١٥ Years Age Group In Baghdad City Khalida Jhalil Ibraheem

Diarrheal disease is defined as having three or more loose or liquid stools per day or more frequently than normal for that person (١). Diarrhea is leading cause of childhood morbidity and mortality in the developing countries , especially in areas with poor disinfection , lake of safe drinking water and inadequate sanitation (٢) . Various etiological agents , including viruses , bacteria , protozoa and helminthes , cause diarrhea . These organisms are transmitted from the stool of one individual to the mouth of another , a route termed feecal – oral transmission . However , they differ in the exact route of entry from stool to mouth and in the infectious dose needed to cause the disease (٣) .

The bacterial species implicated include serotypes of *Escherichia coli* , *Shigella sp.* , *Salmonella sp.* , *Vibrio cholera* , *Compylobacter jejuni* , *Staphylococcus sp.* , *Citrobacter sp.* etc (٤) . *Giardia lamblia* , *Entamoeba histolytica* , *Iodomoeba bullshili* are some of protozoa implicated . Fungal enteritis has also been documented (٥) . Rotavirus is the most common pathogen witch causes gastroenteritis in developing countries (٦) . An epidemiologic study of an infective disease in a community is always considered to be an initial step to the introduction of the appropriate interferences for controlling the disease because the feature and the patterns of isolation of pathological causes of the disease vary from place to place depending on the local geography , metrology and socio – economic elements (٧) .

This study was carried because , diarrhea and associated of droughts state remain the major problem to the public health importance in the medial east especially in the age under ٥ years old . The purpose is to study the various causative factors of diarrheal disease and diagnosis the effect result of it in children under ١٥ years old .

Materials an Methods

A total of ٢٠٠ sample of the stool were collected from children between ٠-١٥ years of age visiting child's central Teaching Hospital and Alkadhmia Hospital for children in the period between April , ٢٠١٦ to October , ٢٠١٦ .

Data collected by mean of study questionnaires . Fresh stool specimens was collected from the patient in to a sterile container and examined microscopically for cysts and trophozoites of parasites , using normal saline and iodine staining wet mounts (٨) .

Standard culture and identification methods were used for bacterial investigation (٩,١٠) . For rotavirus , stool samples were analyzed using

Clinical Study Of Enteropathogenic Which Causes Diarrhea Among ٠-١٥ Years Age Group In Baghdad City Khalida Jhalil Ibraheem

Rota Rapid Diagnostic kit as described by manufacturer (Diaspot company, Indonesia).

Results and Discussion

Out of ٢٠٠ cases in the study group , ٧٢% showed single etiology and ٢٨% mixed etiology Table (١) . Similar observation were made by (١١) . As they found ٢٥% cases of mixed infection . Figure (١) shows that the highest infection rate recorded in the bacterial group (٥١,٥ %) followed by parasitic group (٣٧,٥%) and viral group (١١%) . Similar observation were made by *Saeed et al .*, ٢٠١٥ which found that the prevalence of diarrhea caused by bacteria was significantly higher than that caused by parasitic and virus infection (١٢) .

Distribution of etiological agent according to the age categories shown in Table (٢) , it was observed that maximum bacterial infection rate at age group (< ١ years) were (٤٦%) and Lowest percentage appeared at age group (> ٥ years) were (٢٠%). As for parasitic infections , the most age group susceptible to infection were (> ٥ years) at percentage (٥٢%) and lowest (< ١ years) at percentage (١٩%) . Viral infections recorded the highest infection rate in the age group (١-٥ years) by (٧٣%) and the lowest rate in the age group (> ٥ years) by (٩%) . A similar pattern of age distribution has been found in earlier done studies (١٣,١٤) . Children of less than ٥ years has been identified as the most infection with diarrhea and equally dire consequences. The cause of infection in large age is that these children are more free , active and active in school and outside the home , while not respecting the health conditions (١٥) .

Table (٣) shows the number of days in which the case of diarrhea continued through it is possible to determine whether the infection with diarrhea is acute or chronic . In bacterial infections , most cases were characterized by chronic diarrhea with ٤٩% and ١٩% of children with diarrhea for ١-٢ days . As for parasitic infections , the proportions are almost equal to ٤٠% for continued cases of diarrhea ١-٢ days and ٣٨% more than ٤ days . while most cases of diarrhea for viral infections were acute diarrhea of (٥٩%) and (٣٢%) of children with diarrhea for ١-٢ days .

These results show that most bacterial cases were in the chronic diarrhea stages . Parasitic infections were half of the cases with severe diarrhea and the other half suffered from chronic diarrhea . While viral infections were in the stages of acute diarrhea . These results correspond to a study conducted by (١٦,١٧) .

Clinical Study Of Enteropathogenic Which Causes Diarrhea Among 1-5 Years Age Group In Baghdad City Khalida Jhalil Ibraheem

Distribution of different epidemiological factors are shown by table(٤) . The rate of the diarrhea was higher in male children (٥٨) than in female children (٤٢) , as reported in many previous studies by (١٨,١٩) . In other study by (٢٠) found that male and female children were equally affected . The type of breast feeding was studied in the cases of diarrhea in infants , with the highest incidence of diarrhea in infants who rely on artificial feeding (٧٣%) , while the lowest percentage of infection in infants who rely on breast feeding (٢٧%) . On the basis of drinking water , the highest infections rate among children . Who rely on tap water (٧١,٥%) compared to a sterile water (٢٨,٥%) . No previous results were recorded against this study in terms of reliance on artificial feeding and drinking from tap water which had the highest incidence of diarrhea in children . In a study accompanied by (٢١) , tap water recorded the highest rate of infection with diarrhea (٥١,١%) , as well as the type of artificial feeding (٥٥,٣%) . This study showed that diarrhea is more common in children living in rural setup (٦٠,٥%) than the one living the urban setup(٣٩,٥). This can be ascribed to poor quality of disinfected condition and sanitation practices in rural population contributes high risk of infection especially among children (٢٢).

Table(٥) showed that *E.coli* was recorded as the predominant bacteria with (٢٠,٥) of prevalence followed by *Shigella sp*, *Salmonella sp* and *Klebsiella* with (١٠%), (١٤,٥%) and (٦٠,٥%) respectively. This pattern was similar to other study done by (٢٣) . *Entamoeba histolytica* recorded highest infection rate (١٩,٥%) followed by *G.lambliia* and *B.coli* with (١٢,٥%) and (٥,٥%) respectively. Similar observation were made by (٢٤,٢٥). The percentage of sample infected with the rotavirus was (١١%) compared with the total number of sample studied. This result was similar to the finding of the studies done by (٢٦,٢٧).

Determination of diarrhea etiology and improved hygiene are important for clinical management and controlled strategic planning to reduce the burden of the preventable infectious diseases among children.

Table (١) : Single and mixed pathogens identified in stool sample from diarrheal cases .

Etiology	No. of cases	Percentage (%)
Single pathogen	١٤٤	٧٢
Mixed pathogen	٥٦	٢٨
Total	٢٠٠	١٠٠

**Clinical Study Of Enteropathogenic Which Causes Diarrhea Among
 ٠-١٥ Years Age Group In Baghdad City Khalida Jhalil
 Ibraheem**

Table (٢) : The percentage of etiological agent according to the age Categories .

Etiology	< ١ years(%)	١- ٥years(%)	>٥ years(%)	Total(%)
Bacteria	٤٧(٤٦)	٣٥(٣٤)	٢١(٢٠)	١٠٣(١٠٠)
parasite	١٤(١٩)	٢٢(٢٩)	٣٩(٥٢)	٧٥(١٠٠)
Virus	٤(١٨)	١٦(٧٣)	٢(٩)	٢٢(١٠٠)
Total	٦٥(٣٢,٥)	٧٤(٣٧)	٦١(٣٠,٥)	٢٠٠(١٠٠)

Table (٣) : The percentage of the etiological agent according to the duration of the time for continuation of diarrheal cases .

Etiology	١-٢Day(%)	٣-٤Days(%)	>٤Days(%)	Total(%)
Bacteria	٢٠(١٩)	٣٣(٣٢)	٥٠(٤٩)	١٠٣(١٠٠)
Parasite	٣٠(٤٠)	١٧(٢٣)	٢٨(٣٧)	٧٥(١٠٠)
Rotavirus	١٣(٥٩)	٧(٣٢)	٢(٩)	٢٢(١٠٠)
Total	٦٣(٣١,٥)	٥٧(٢٨,٥)	٨٠(٤٠)	٢٠٠(١٠٠)

Table(٤): Distribution of different epidemiological factors among the children offered with diarrhea.


Epidemiological Factors		N=٢٠٠ No(%)
Sex	Male	١١٦(٥٨)
	Female	٨٤(٤٢)
Feeding	Artificial Feeding	١٤٦(٧٣)
	Breast Feeding	٥٤(٢٧)
Water source	Tap Water	١٤٣(٧١,٥)
	Sterile Water	٥٧(٢٨,٥)
Residence	Rural	١٢١(٦٠,٥)
	Urban	٧٩(٣٩,٥)

Table(٥): Prevalence of various etiological agent in diarrhea patients.

Etiology	Number(%)
Bacteria	
<i>Escherichia coli</i>	٤١(٢٠,٥)
<i>Salmonella sp.</i>	٢٠(١٠)
<i>Shigella sp.</i>	٢٩(١٤,٥)
<i>Klebsiella sp.</i>	١٣(٦,٥)
Parasite	

**Clinical Study Of Enteropathogenic Which Causes Diarrhea Among
 ٠-١٥ Years Age Group In Baghdad City Khalida Jhalil
 Ibraheem**

<i>Entamoeba histolytica</i>	٣٩(١٩,٥)
<i>Giardia lamblia</i>	٢٥(١٢,٥)
<i>Blantidium coli</i>	١١(٥,٥)
Virus	
Rotavirus	٢٢(١١)
Total	٢٠٠(١٠٠)



Figure(١): The percentage of the etiological agents causing diarrhea .

References

- ١- Kosek M. ; Bern C. and Guerrant R.L.(٢٠٠٣) . The global burden of diarrhoeal disease as estimated from studies published between ١٩٩٢ and ٢٠٠٠ . Bull World Health Organ . ٨١ : ١٩٧ – ٢٠٤ .
- ٢- Bruce J.; Boschi-Pinto C.; Shibuya K.and Back R.E.(٢٠٠٥) . WHO estimates of the causes of death in children . Lancet. ٣٦٥ : ١١٤٧-١١٥٢.
- ٣- Kerri B.; Christine M.; Khavari N.and Manji K. (٢٠١٧) .Etiology of diarrhea , nutritional outcomes , and novel intestinal biomarkers in tanzanian infants . J Pedia Gastro & Nutri . ٧٦٤ –Issue ١ : ١٠٤-١٠٨ .

**Clinical Study Of Enteropathogenic Which Causes Diarrhea Among
٠-١٥ Years Age Group In Baghdad City Khalida Jhalil
Ibraheem**

- ٤- Rangel –Vargas E. ; Gomes-Aldapa C.A.; Torres-Vitela Mdel R.; Villarruel-Lopez A.; Gordillo-Martinez A.J.and Castro-Rosas J.(٢٠١٥). Presences and correlation of some enteric indicator bacteria, diarrheagenic *Escherichia coli* pathotypes, and *Salmonella* serotypes in alfalfa sprouts from local retail markets in Pachuca, Mexico. J Food Prot ٧٨(٣):٦٠٩-٦١٤.
- ٥- Moreno A.C.; Filho A.F.; Gomes Tdo A. ; Ramos S.T.; Montermor L.P.and Tavares V.C.(٢٠١٠). Etiology of childhood diarrhea in the northeast of Brazil : Significant emergent diarrheal pathogens .Diagn Microbiol Infect Dis .٦٦: ٥٠-٥٧.
- ٦- Bonkougou I.J.; Sanou I.; Bon F.; Benon B.; Coulibaly S.O.; Haukka K.; Traore A.S.and Barro N.(٢٠١٠). Epidemiology of rotavirus infection among young children with acute diarrhea in Burkina Faso . BMC Pediatr . ١٠:١٠-٩٤.
- ٧- Guarino A.; Ashkenazi S.; Gendrel D.; (٢٠١٤) . European Society for pediatric Gastroenterology , Hepatology , and Nutrition /European Society for Pediatric Infectious Diseases evidence-based guidelines for the management of acute gastroenteritis in children in Europe. J Pediatr Gastroenterol Nutr . ٥٩ :١٣٢.
- ٨- Melvin, D. M.; and Healy, G. R .(١٩٨٥). Intestinal and urogenital protozoa. In maunual of clinical microbiology (٤th ed) .Lennette .E . H.; Balows, A.; Hausler ,W .J .; and Shadomy ,H .J .Eds .American Society for Microbiology ,Washington .
- ٩- Murray P. R.; Baron E. J.; Jorgensen R.H.; Pfaller M. A. and Yollken R.H.(٢٠٠٣). Manual of clinical microbiology , ٨th ed .ASM Press , Washonton , DC.
- ١٠- Gillespie S.H.and Hawkey P.M.(٢٠٠٦). Principles and Practice of Clinical Bacteriology .Chischester : John Wiley & Sons, ٢ .
- ١١- Kumar M.; Ramkrishana M.and Kumar L.(٢٠١٤). Study of various causative factors of diarrhoeal diseases with special reference to bacteriological profile among ٠ to ٥ years age group . I.J.S.S. Vol ٢.Issue ٣.
- ١٢- Saaed A.; Abd H.and Sandstrom G.(٢٠١٥). Microbial etiology of acute diarroea in children under five years of age in Khartoum , Sudan.J Med Microbiol. ٦٤(pt ٤):٤٣٢-٤٣٧ .
- ١٣- Poo M.I. and Lee W.S (٢٠٠٧). Admission to hospital with childhood acute gastroenteritis in Kuala Lumpur , Malaysia . Med J Malaysia . ٦٢(٣) : ١٨٩-٩٣.
- ١٤- Yongsi H.B.N.(٢٠٠٨). Pathogenic microorganisms associated with childhood diarrhoeain low and middle income countries . Int J. Environ Res Public Health. ٥:٢١٣-٢٩ .

**Clinical Study Of Enteropathogenic Which Causes Diarrhea Among
٠-١٥ Years Age Group In Baghdad City Khalida Jhalil
Ibraheem**

- ١٥- Anasari S.; Bahadur S.; Parajuli K.; Paudyal R.P.A. (٢٠١٢). Pattern of acute parasitic diarrhea in children under five years of age in Kathmandu, Nepal. Open Journal of Medical Microbiology ,٢:٩٥-١٠٠.
- ١٦- Wang,Y.; Dennehy, P.H.; Keyserling, H.L.; Tang, K. (٢٠٠٧). Rotavirus Infection Alters Peripheral T-Cell Homeostasis in Children with Acute Diarrhea. Journal of Virology, ٨١(٨): ٣٩٠٤-٣٩١٢.
- ١٧- Noaman, A.A. (٢٠١٣). Acute and chronic diarrhea among hospitalized infants aged (١-١٢) months . Diyala Journal for Pure Sciences, ١٠(١): ١-١٠.
- ١٨- Sherchand J.B.; Yokoo M.; Sherchand O.; Pant A.R. and Nakogomi O. (٢٠٠٩). Burden of enteropathogens associated diarrheal disease in children hospital, Nepal .Sci World .٧,٧١-٧٥.
- ١٩- Moyo S.J.; Gro N.; Mateel M.I. and Kilundu J. (٢٠١١). Age specific aetiological agents of diarrhea in hospitalized children age less than five years in Dar es Salaam ,Tanzania. BMC Pediatrics ١١:١٩-٢٤.
- ٢٠- Mashoto K.O.; Malebo H.M.; Msisiri E. and Peter E. (٢٠١٤). Prevalence , one week incidence and knowledge on causes of diarrhea : household survey of under- five and adults in Mkurange distric, Tanzania . BMC Public Health ١٤,٩٨٥,١٠.
- ٢١- Hasan, A.Sh.; Al-Duliami, A.A.; Abdul-Kadir, M.Sh. and Al-Azawi, Y. (٢٠١١). The prevalence of Rotavirus Infection in Baquba - Diyala province. Iraqi J. Comm. Med., July., ٢٤(٣): ٢٠٠-٢٠٤.
- ٢٢- Diniz-Santos D.R.; Santana J.S.; Barreto J.R.; Andrade M.G.M. and Silva L.R. (٢٠٠٥). Epidemiological and microbiological aspects of acute bacterial diarrhea in children from Salvador, Bahia, Brazil. Braz J Infect Dis . ٩(١) .
- ٢٣- Subashkumar R.; Thayumanavan T.; Vivekanandhan G. and Lakshmanaperumalsay P. (٢٠١٢). Etiology of childrens diarrhea in Southern India: Associated pathogens and usual isolates . Afr .J. Microbiol. Res. ٦(١١): ٢٨٠٨-١٥.
- ٢٤- Shrestha S.K.; Rai S.K.; Vitrakoti R. and Pokharel P. (٢٠٠٩). Parasitic infection in school children in Thimi area, Kathmandu Valley . J Nep Assoc Med Lab Sci ١٠,٣١-٣٣.
- ٢٥- Thapa Magar D.; Rai S.K.; Lekhak B. and Rai K.R. (٢٠١١). Study of parasitic infection among children of Sukumbasi Basti in Kathmandu valley . Nepal Med Coll J ١٣,٧-١٠.
- ٢٦- Nitiema, L.W.; Nordgren, J.; Ouermi, D.; Dianou, D. (٢٠١٠). Burden of rotavirus and other enteropathogens among children with diarrhea in Burkina Faso. Int. J. Infect. Dis., ١٥(٩): ٤٦-٥٢.
- ٢٧- Al-Ameen, H.A.; Al-Hmudi , H.A. and Darush, A.A. (٢٠١٢). Epidemiology of Rotavirus cases among children under age ٥ years in Basra province from ٢٠٠٨-٢٠١١. J.Thi-Qar Sci., ٣(٣): ٣١-٣٦.

الخلاصة

Clinical Study Of Enteropathogenic Which Causes Diarrhea Among ٠-١٥ Years Age Group In Baghdad City Khalida Jhalil Ibraheem

تعد امراض الاسهال سببا رئيسيا للوفيات بين الاطفال في البلدان النامية. صممت هذه الدراسة لتحديد المسببات البكتيرية والفيروسية والطفيلية والعوامل السريرية والوبائية لدى الاطفال الذين تتراوح اعمارهم بين ٠ - ١٥ سنة والذين يعانون من الاسهال في مدينة بغداد. تم جمع ٢٠٠ عينة براز من الاطفال المصابين بالاسهال خلال مراجعتهم مستشفى الطفل المركزي التعليمي ومستشفى الكاظمية للأطفال للفترة ما بين نيسان ٢٠١٦ الى تشرين الاول ٢٠١٦. شخضت البكتريا المعزولة باستخدام الطرق التشخيصية والزرعية القياسية . وبالفحص المجهرى تم تشخيص الطفيليات باستخدام المحلول الملحي الفسيولوجي والشرائح الرطبة المصبغة باليود اما الفيروس العجلي الدوار فتم الكشف عنه بواسطة عدة التشخيص السريع. من مجموع ٢٠٠ عينة براز شخضت ١٤٤ عينة مخمجة بنوع واحد من المسببات المرضية بنسبة ٧٢% و ٥٦ عينة تحوي اخماج مشتركة بنسبة ٢٨% . ان اعلى نسبة خمج كانت للمجموعة البكتيرية (٥١,٥%) تليها المجموعة الطفيلية (٣٧,٥%) والمجموعة الفيروسية (١١%). وفقا للفئات العمرية المدروسة لوحظ ان اعلى نسبة للاخماج البكتيرية كانت عند الفئات العمرية اقل من سنة واحدة بنسبة (٤٦%) بينما سجلت الاخماج الطفيلية اعلى نسبة خمج عند الفئة العمرية اكثر من ٥ سنوات بنسبة (٥٢%) اما الاخماج الفيروسية فقد كانت الفئة العمرية ١-٥ سنوات اكثر عرضة للخمج بنسبة (٧٣%) . اظهرت نتائج الدراسة ان البكتريا السائدة هي *Escherichia coli* بنسبة ٢٠,٥% تليها *Shigella sp.* , *Salmonella sp.* و *Klebsiella sp.* بنسبة ١٤,٥% ، ١٠% ، ٦,٥% على التوالي . سجلت *Entamoeba histolytica* اعلى معدل للاخماج الطفيلية (١٩,٥%) تليها *Giardia lamblia* (١٢,٥%) و *Blantidium coli* (٥,٥%) وكانت النسبة المئوية للعينات المصابة بالفايروس العجلي الدوار ١١% مقارنة مع مجموع العينات التي تمت دراستها .